Referral Status: 🛛 New Patient 🎵 Updated Patient Information



CMC Infusion Center- Referral Face Sheet

Patient Name:	Patient D.O.B	
Current Height:	Current Weight	
Patient Phone # ()	Gender	
Patient Address		
CityState	_Zip Code	
Related Diagnosis Code (ICD-10 code):		
Referring Provider (Please Print)		
Office Phone # ()	Office Contact:	

If referring provider and/or patient is external to Conway Medical Center. Please include:

- Most recent history and physical including:
 - \circ Comprehensive medication list
 - o Past medical history
 - \circ Related past/ failed therapies (with dates)
 - Known allergies
- Most recent labs / test result (Ex. CBC, PPD, Hepatitis screening, etc.)
- Insurance Information

Please fax requested documentation, face sheet, and treatment plan to **843-234-5460** attention CMC Medication Management Specialist. Please feel free to call 843-234-8575 with any questions.

We look forward to the opportunity to be a part of your patient's treatment journey at Conway Medical Center.

Referral Status 🛛 New 🖾 Order Change 🗖 Order Renewal

IVIG Treatment Plan



Patient Name:	Patient DOB	
General Diagnosis-	ICD-10 Code	
Pt. Weight Pt He	ight Known Allergies	
Requested Start Date//		
Has patient previously received th	is medication \square No \square Yes, if so, date of last infusion	י/
Authorized Treatment Duration	□ Infuse Once □ Administer every days for _	doses
Infusion (Month 1 to 12) Durat	ion: 12 Months	
Vital Signs (Month 1 to 12 every 30 min, Resp to discharge) ɔ, BP, HR, Temperature on admission to the unit and prior	to infusion HR, BP and prior
Pharmacy may substitute IV	/IG product based on insurance approval (Gamunex, Octa	gam, Privigen, etc.)
Provider requests specific IV	VIG product:	
Pre- Treatment Medications Tylenol (Month 1 to 12)		
	. Tab, Day of Tx t s: Administer 30 minutes before dose	
25 mg, IV Pusl	h, Injection, Day of Tx t s: Administer 30 minutes before dose	
50 mg, IV Push Comment SOLU-Medrol (Month 1 to 12)	h, Injection, Day of Tx t s: Administer 30 minutes before dose	
40 mg, IV Push	h, Powder-Inj, Day of Tx t s: Administer 30 minutes before dose	
L 125 mg, IV Pus Comment	h, Powder-Inj, Day of Tx t s: Administer 30 minutes before dose	
,	onse to any live vaccines. Such vaccines should be avoided	

administration: Measles, and Varicella immunizations should be deferred for greater than or equal to 11 months after receiving IVIG. IVIG should be administered at the minimum dose and rate possible in patients at risk of renal dysfunction. Referring providers agrees to monitor renal function and notify Conway Infusion Services if renal function deteriorates resulting in therapy discontinuation. IVIG has been associated with risks of thrombotic events and should be administered at the minimum dose and infusion rate possible.

Prescriber Signature (No Stamped Signatures or Electronic Signatures)

Provider S	ignatı	ıre		Provider Name (please print)_	
Date	/	_/	Patient Name	Ра	atient DOB

IVIG Treatment Plan

			Provider Initials
IVIG	_		
		1 g/kg, IV Piggyback, Injection, Once Comments: Initial rate = 0.5 mg/kg/m to 5 mg/kg/minute (6 mL/kg/hour)	inute (0.6 mL/kg/hour); May Increase slowly (if tolerated) up
		1 g/kg, IV Piggyback, Injection, every 24 Comments: Initial rate = 0.5 mg/kg/m to 5 mg/kg/minute (6 mL/kg/hour)	hr for doses inute (0.6 mL/kg/hour); May Increase slowly (if tolerated) up
		400 mg/kg, IV Piggyback, Injection, ever Comments: Initial rate = 0.5 mg/kg/n to 5 mg/kg/minute (6 mL/kg/hour)	y 24 hr fordoses hinute (0.6 mL/kg/hour); May Increase slowly (if tolerated) up
		500 mg/kg, IV Piggyback, Injection, Onc Comments: Initial rate = 0.5 mg/kg/m to 5 mg/kg/minute (6 mL/kg/hour)	e ninute (0.6 mL/kg/hour); May Increase slowly (if tolerated) up
		Other:mg infused every 24 h Comments: Initial rate = 0.5 mg/kg/r up to 5 mg/kg/minute (6 mL/kg/hour,	ninute (0.6 mL/kg/hour); May Increase slowly (if tolerated)
Commur	Cor		s, rigors, hypotension or respiratory difficulty during IVIG
Commur			
	Verify		n the last 6 months Measles and Varicella immunizations 1 months after receiving IVIG.
Notify Pr	Not	(Month 1 to 12) fy provider for temperature > 99.5 or HR seline	< 55 or > 140, drop or rise in SBP more than 20 mm Hg from
Notify Pr	Not	(Month 1 to 12) fy provider for any live vaccine administra arance is received.	tion within the last 6 months and hold IVIG until provider
INF INF	nfusio	n Room Orders Subphase (Month 1 to 12)
	o the r Pre	n Disclaimer : Per CMC P&T approval, IV earest manufacturer vial size by pharmac gnant patients: Dose using actual body w n-pregnant patients:	
		 Use actual body weight if actual bod (IBW). 	y weight (ABW) is less than 120% of ideal body weight
	o Ma	 Ose ADJOSTED body weight if actu weight (IBW). les: IBW = 50 kg + 2.3 kg for each inch or 	al body weight (ABW) is greater than 120% of ideal body /er 5 ft
•		nales: IBW = 45.5 kg +2.3 kg for each inc	
Prescriber	Sign	ature (No Stamped Signatures or E	lectronic Signatures)
Provider Sig	gnatu	re Provi	der Name (please print)
Date/		Patient Name	Patient DOB

IVIG Treatment Plan

• ADJUSTED body weight = IBW + 0.4(ABW – IBW)

Prescriber Signature (No Stamped Signatures or Electronic Signatures)

Provider Signature _		_ Provider Name (please print)
Date//	Patient Name	Patient DOB